

Phys B - week 4 - Giancoli Ch 9 # 5, 11, 53, 78, 89

Giancoli Ch 10 # 6, 17, 53, 57, 86

5. (II) The force on a particle of mass  $m$  is given by  $\mathbf{F} = 26\mathbf{i} - 12t^2\mathbf{j}$  where  $F$  is in N and  $t$  in s. What will be the change in the particle's momentum between  $t = 1.0\text{ s}$  and  $t = 2.0\text{ s}$ ?

$$\vec{F} = F_x \mathbf{i} + F_y \mathbf{j}$$

$$F_x = 26 \quad F_y = -12t^2$$

$$\vec{F} = \frac{d\vec{p}}{dt} \quad F_x = \frac{dp_x}{dt} = 26 \quad F_y = \frac{dp_y}{dt} = -12t^2$$

$$\vec{p} = \int \vec{F} dt \quad p_x^{(t)} = \int F_x dt = \int 26 dt = 26t$$

$$p_y(t) = \int_{t=1}^t F_y dt = \int -12t^2 dt = -\frac{12}{3} t^3 = -4t^3$$

$$\Delta p_x = 26t \Big|_{t=1}^{t=2} = 26(2-1) = 26 \text{ kg m/s}^2$$


$$\Delta p_y = -4t^3 \Big|_{t=1}^{t=2} = -4(2^3 - 1^3) = -4(8-1) = -4(7) = -28 \text{ kg m/s}^2$$

*Escherichia coli*

$$f_a = \frac{dV}{dt}$$

Cu. 9 # 11. (Matt E + Ingrid L)

11. (I) An atomic nucleus at rest decays radioactively into an alpha particle and a smaller nucleus. What will be the speed of this recoiling nucleus if the speed of the alpha particle is  $2.5 \times 10^5 \text{ m/s}$ ? Assume the nucleus has a mass 57 times greater than that of the alpha particle.



$$\sum \vec{F} = 0 = m_n V_n - m_\alpha V_\alpha$$

$$m_n V_n = m_\alpha V_\alpha$$

$$\text{Find } V_n = \frac{m_\alpha V_\alpha}{m_n}$$

$$m_N = 57 m_\alpha$$

$$V_\alpha = 2.5 \cdot 10^5 \frac{\text{m}}{\text{s}} = \\ 57$$

$\partial h / \partial s$

$$M_F \\ + h)^2$$

$$\frac{dV}{dt}$$


$y_p$

BEFORE

53. (II) An atomic nucleus of mass  $m$  traveling with speed  $v$  collides elastically with a target particle of mass  $2m$  (initially at rest) and is scattered at  $90^\circ$ . (a) At what angle does the target particle move after the collision? (b) What are the final speeds of the two particles? (c) What fraction of the initial kinetic energy is transferred to the target particle?


Find  $\theta$

Find  $v_1, v_2$



Solve for  $\theta, v_1, v_2$

in terms of  $v, m$



Initially (BEFORE)  $\rightarrow P = mv$  in the  $x$ -dir

AFTER: in the  $x$ -direction

$$P_{2x} =$$

$$\text{Since } p_1 = p_{1x} = p_{2x}$$

(1)

$$\text{BEFORE: } p_y = 0$$

$$P_{2y}$$

$$\text{AFTER: } p_{2y} =$$

$$p_{1y} =$$

$$p_y \text{ before} = p_y \text{ after}$$


(2)

We have 2 eggs in 3 vulcanoes: ①  
\_\_\_\_\_ ②

We need a third equation: ENERGY CONSERVATION  
works for elastic collisions

BEFORE

AFZER



$$\frac{1}{2}mV^2 = \frac{1}{2}mv_1^2 + \frac{1}{2}(2m)v_2^2$$

3

Now solve 3 equations in 3 unknowns

Hint - Square and add ① and ② first:


$$(Jada) \cdot h \cdot 10^6 \text{ m} = R_E \text{ by coincidence!}$$

- \*78. (II) A rocket traveling 1850 m/s away from the Earth at an altitude of 6400 km fires its rockets, which eject gas at a speed of 1200 m/s (relative to the rocket). If the mass of the rocket at this moment is 25,000 kg and an acceleration of  $1.7 \text{ m/s}^2$  is desired, at what rate must the gases be ejected?

$v_{\text{gas}}$


$$a = \frac{dv_r}{dt}$$

$M$



$$\uparrow a = \frac{dv}{dt}$$

Earth's rest frame: Rocket's rest frame:



$h$

Gravity @  $h$  is

$$g' = G \frac{M_E}{(R_E + h)^2}$$


$\downarrow R_{\text{Earth}}$

Actually, we are told that  $v_{\text{gas}} = 1200 \frac{\text{m}}{\text{s}}$  relative to rocket

Momentum conservation of

rocket-gas system:

$$\downarrow p_{\text{gas}} = p_{\text{rocket}} \uparrow$$



$$v_{\text{gas}} \frac{dm}{dt} = F_{\text{up}}$$

$$\uparrow a = \frac{dv}{dt}$$


Forces on rocket: Gravity ↓ and Thrust ↑

$$\Sigma F = Ma$$

$$F_{\text{up}} - Mg' = Ma = v_{\text{gas}} \frac{dm}{dt} - Mg'$$

$$\text{Gas must be ejected at rate } \frac{dm}{dt} = M(a + g') / v_{\text{gas}}$$

99. The gravitational slingshot effect in Fig. 9-52, shows the planet Saturn moving in the negative  $x$  direction at its orbital speed (with respect to the Sun) of 9.6 km/s. The mass of Saturn is  $5.69 \times 10^{26}$  kg. A spacecraft with mass 825 kg approaches Saturn, moving initially in the  $+x$  direction at 10.4 km/s. The gravitational attraction of Saturn (a conservative force) causes the spacecraft to swing around it (orbit shown as dashed line) and head off in the opposite direction. Estimate the final speed of the spacecraft after it is far enough away to be nearly free of Saturn's gravitational pull.




We can neglect the change in  
Saturn's speed due to the  
momentum transfer, since it is so

$$\text{massive: } dV_{\text{spaceship}} / m_{\text{spaceship}} = dV_{\text{Sat}} / m_{\text{Sat}}$$

$$m_{\text{Saturn}} = m_{\text{spaceship}} \approx 10^{26} \text{ kg}^{-24}$$

$$dV_{\text{spaceship}} \quad m_{\text{saturn}} \quad 10^{26} \text{ kg}^{-10}$$


Transform to a new frame




Grancoli Ch 10 #6, 17, 53, 57, 86

6. (II) Calculate the angular velocity of the Earth (a) in its orbit around the Sun and (b) about its axis.

$$(a) L = m v d = \frac{m d}{T} (2\pi r)$$



$$L_{\text{orbit}} = \frac{2\pi m d^2}{T}$$



$$L_{\text{spin}} = I \omega \text{ where } \omega = \frac{2\pi \text{ rad}}{1 \text{ day} \cdot 24 \text{ h} / 3.6 \times 10^3 \text{ s}}$$

$$\omega_{\text{spin}} = \frac{\text{rad}}{\text{s}}$$

$$\text{Moment of inertia of sphere } I = \frac{2}{5} M R^2$$

Assume Earth has uniform density:

$$I = \frac{2}{5} (\rho)(M)(R)^2 = \frac{\rho M R^2}{5}$$



$$L_{\text{spin}} = \left( \frac{\text{rad}}{\text{s}} \right) \left( \frac{\text{kg m}^2}{\text{s}} \right) = \frac{\text{kg m}^2}{\text{s}}$$

57. (II) Determine the angular momentum of the Earth (a) about its rotation axis (assume the Earth is a uniform sphere), and (b) in its orbit around the Sun (treat the Earth as a particle orbiting the Sun). The Earth has mass =  $6.0 \times 10^{24} \text{ kg}$ , radius =  $6.4 \times 10^6 \text{ m}$ , and is  $1.5 \times 10^9 \text{ km}$  from the Sun.


Oops - I answered this above, Still need

$$(a) L_{\text{orbit}} = \frac{2\pi \text{ rad}}{1 \text{ yr} \cdot \pi \times 10^7 \text{ s}}$$

In Quantum Mechanics, we will similarly consider the spin and orbital angular momentum of particles, e.g. of an electron in orbit around a proton



17. (II) The angular acceleration of a wheel, as a function of time, is  $\alpha = 5.0t^2 - 3.5t$ , where  $\alpha$  is in rad/s<sup>2</sup> and  $t$  in seconds. If the wheel starts from rest ( $\theta = \omega = 0$  at  $t = 0$ ), determine a formula for (a) the angular velocity  $\omega$  and (b) the angular position  $\theta$ , both as a function of time. (c) Evaluate  $\omega$  and  $\theta$  at  $t = 2.0\text{ s}$ .



$$s = r\theta$$

$$v = r\omega$$

$$\omega = \frac{d\theta}{dt}$$

$$\alpha = r\ddot{\theta} \quad \ddot{\theta} = \frac{d\omega}{dt}$$

Since  $\alpha = \frac{d\omega}{dt}$ , then  $\omega = \int \alpha dt$

(a)  $\omega = \int (5t^2 - \frac{7}{2}t) dt =$

(b)  $\theta = \int \omega dt =$

(c)  $\omega(2\text{s}) =$

$\theta(2\text{s}) =$

53. (I) A diver (such as the one in Fig. 10-30) can reduce her moment of inertia by a factor of about 3.5 when changing from the straight position to the tuck position. If she makes two rotations in 1.5 s when in the tuck position, what is her angular speed (rev/s) when in the straight position?

STRAIGHT DIVE      TUCK DIVE  
 $I_0$        $I = I_0 = I_0 = 2.0 = 2.20$   
 slower      faster       $\frac{3.5}{7/2} = \frac{7}{2} = 7$

Angular momentum is conserved (if external torques = 0)

$$P = \mu v$$

$$L_{\text{straight}} = L_{\text{tuck}}$$

$$L = I\omega$$

$$I_0 \omega_0 = I_T \omega_T$$

$$\omega_0 = \omega_T \frac{I_T}{I_0} = \omega_T \frac{2}{7}$$

$$\omega \left( \frac{\text{rad}}{\text{s}} \right) = 2\pi f \left( \frac{\text{rev}}{\text{s}} \right)$$


$$\text{Angular speed in tuck is } f_T = \frac{2 \text{ rev}}{3/2 \text{ sec}} = \frac{\text{rev}}{\text{sec}}$$

$$\text{Angular speed in straight dive is } f_0 = f_T \frac{2}{7} = \frac{\text{rev}}{\text{sec}}$$

Greater  $I_0 \rightarrow$  Slower  $f_0$

86. The Moon orbits the Earth so that the same side always faces the Earth. Determine the ratio of its spin angular momentum (about its own axis) to its orbital angular momentum. (In the latter case, treat the Moon as a particle orbiting the Earth.)

Looking down at N pole: ☽



Orbital angular momentum

$$L_o = m V R_o = m R_o \left( \frac{2\pi R_o}{T_o} \right)$$

$$T_o \approx 28 \text{ d} \left| \frac{3.6 \cdot 10^3 \text{ s}}{1 \text{ d}} \right| = \text{ s}$$


$$L_o = \frac{2\pi m R_o^2}{T_o}$$

orbital period ~ 4 weeks

Spin angular momentum

$$p = m v$$

$$L_s = I w_s$$



Assume moon is a constant-density sphere;  $I = \frac{2}{5} M R_m^2$

Phase locked: we always see the same face of the moon because its spin period  $T_s = T_o$  orbital period,  $w_s = \frac{2\pi}{T_s}$

$$L_s = \frac{2}{5} M R_m^2 \frac{2\pi}{T_o}$$

$$\text{So } \frac{L_s}{L_o} = \frac{\frac{2}{5} 2\pi M R_m^2 / T_o}{2\pi m R_o^2 / T_o} = \frac{2(R_m)^2}{5(R_o)^2} = \frac{2}{5} \left( \frac{R_m}{R_o} \right)^2$$

$$\frac{L_s}{L_o} =$$